

The Great Basin Naturalist

PUBLISHED AT PROVO, UTAH, BY
BRIGHAM YOUNG UNIVERSITY

ISSN 0017-3614

VOLUME 55

31 JULY 1995

No. 3

Great Basin Naturalist 55(3), © 1995, pp. 193–200

BENTHIC COMMUNITY STRUCTURE IN TWO ADJACENT STREAMS IN YELLOWSTONE NATIONAL PARK FIVE YEARS AFTER THE 1988 WILDFIRES

G. Wayne Minshall¹, Christopher T. Robinson¹, Todd V. Royer¹, and Samuel R. Rushforth²

ABSTRACT.—Physical characteristics, benthic macroinvertebrates, and periphyton assemblages in two adjacent headwater streams in Yellowstone National Park were evaluated five years after the 1988 wildfires. The catchment of one stream was burned by wildfire (burned stream) while the other catchment was unburned (unburned stream). Physical measures revealed channel alteration in the burned stream relative to the unburned stream. Periphyton biomass was lower in the burned than the unburned stream (29.2 vs. 50.5 g/m² AFDM, respectively), further demonstrating the unstable physical conditions of that system. Kendall's coefficient of concordance (an index of similarity) between diatom assemblages was 0.22, indicating distinct assemblage composition between streams. *Navicula permissis* Hust. was the most abundant diatom in the burned stream while *Hannaea arcus* (Ehr.) Patr. was dominant in the unburned stream. Macroinvertebrate taxa richness, density, and biomass were all greater in the unburned stream, although Chironomidae was the most abundant taxon in both streams. Results suggest the removal of terrestrial/riparian vegetation by wildfire can directly influence stream benthic assemblages by altering the inherent disturbance regime of the physical habitat template.

Key words: wildfire, streams, disturbance, macroinvertebrates, diatoms, benthic habitat, Yellowstone National Park.

Physical disturbance, acting at various spatial and temporal scales, often is the predominant factor structuring stream benthic communities (Minshall 1988, Resh et al. 1988). Further, physical disturbances may be viewed in a hierarchical framework, with the effects of small-scale disturbances altered (intensified or mediated) by large-scale disturbance events (sensu O'Neill et al. 1986). Wildfire, as a large-scale disturbance, directly influences stream biotic structure and function by affecting the physical habitat of stream ecosystems (Minshall et al. 1989, Minshall and Brock 1991, Richards

and Minshall 1992, Robinson et al. 1994, Mihuc et al. in press, Robinson and Minshall in press). In lotic ecosystems, physical disturbance also may constrain the establishment of biotic controls, such as competition and predation, on benthic community structure (McAuliffe 1984, Minshall and Petersen 1985, Resh et al. 1988).

Wildfire burned extensive portions of the Greater Yellowstone Ecosystem during the summer of 1988. Over 32% of the streams in Yellowstone National Park (YNP) were affected to varying degrees by wildfires (Minshall et al. 1989, Minshall and Brock 1991, Robinson et al.

¹Stream Ecology Center, Department of Biological Sciences, Idaho State University, Pocatello, ID 83209.

²Department of Botany and Range Science, Brigham Young University, Provo, UT 84602.

1994). Minshall and Brock (1991) summarized the immediate effects of the fires on YNP's stream ecosystems and hypothesized on the mid-term (10–25 yr) and long-term (50–300 yr) effects. They suggested that most adverse short-term effects on streams resulted from increased sediment load and channel erosion caused by increased overland runoff following precipitation events and snowmelt. The intensity and frequency of short-term effects were hypothesized to decrease by year 5 as riparian conditions improve (see Richards and Minshall 1992). In general, mid- and long-term effects on streams, including recovery to prefire conditions, should correspond to vegetative regrowth in burned catchments (Minshall et al. 1989, Minshall and Brock 1991).

The present study compared physical characteristics and benthic community structure in two streams five years after the 1988 wildfires. The streams are adjacent second-order (after Strahler 1952) tributaries of the South Fork Cache Creek. The catchment of one stream was burned during the 1988 wildfires, while the catchment of the other was essentially unburned. The spatial arrangement of these streams (adjacent basins) provided a treatment/reference situation where confounding factors of climate and geology are minimized when comparing differences among the study streams. However, the study lacks true replication of the burned and unburned treatments (*sensu* Hurlbert 1984) and must be viewed as a simple comparison study. Phenomenological studies and/or two stream comparisons are common in stream ecology (e.g., Wallace et al. 1986, Robinson et al. 1993, Scarsbrook and Townsend 1993) and are capable of providing valuable insights (Townsend 1989). The present study can be viewed as a natural "experiment" with observed differences between the two streams attributed to the effects of wildfire. In that context, the study provides insights on general patterns of lotic ecosystem recovery to an unpredictable, large-scale disturbance (Townsend 1989, Lamberti et al. 1991).

METHODS

The study streams, located in the northeast corner of YNP, were surveyed on 19 July 1993. One stream had over 80% of its catchment burned during the 1988 Yellowstone wildfire (hereafter, burned stream; 110°01'30"W,

44°50'00"N), while less than 10% of the catchment of the other stream was burned (hereafter, unburned stream; 110°01'00"W, 44°49'30"N). Climate of the area is typical of the northern Rocky Mountains, with precipitation primarily occurring as snow during the winter months. Both streams drain catchments primarily vegetated (prior to the fire in the burned stream) by coniferous forests of lodgepole pine (*Pinus contorta*) and Engelmann spruce (*Picea engelmannii*). Riparian vegetation consisted of willow (*Salix*), rose (*Rosa*), and alder (*Alnus*).

Surveys were conducted approximately 0.5 km above the confluence of the two streams. Physical characteristics were measured in each stream at five cross-sectional transects, each situated approximately 50 m apart. Measurements made at each transect included stream width at baseflow, stream width at bankfull discharge, and stream cross-sectional profile (for calculation of width:depth ratios). Discharge was calculated in each stream at the most suitable transect following the methods of Platts et al. (1983). In addition to measurements at each transect, 100 randomly selected rock substrata along a 100-m length of stream (located within the outermost cross-sectional transects) were measured for size (length of the longest axis) and percent embeddedness. Embeddedness was defined as the percent coverage of the rock (three-dimensional surface) by fine sediments. Large boulders that protruded through the water surface were not used in substratum size measurements. Water depth and near-bed water velocity also were recorded at each of the 100 random locations. Near-bed water velocity was measured with a small Ott C-1 current meter approximately 2 cm above each substratum.

One periphyton sample was collected from a suitable (flat-surfaced, medium-sized) rock substratum at each cross-sectional transect using a method described in Robinson and Minshall (1986). Samples were frozen in the field in a Taylor-Wharton 3DS dry shipper charged with liquid nitrogen and returned to the laboratory for processing. In the laboratory, samples were extracted in 10 ml of methanol for 24 h (Holm-Hansen and Riemann 1978). One 3-ml subsample was then removed from each sample and analyzed for chlorophyll *a* using a Gilford Instruments (Model 2600) spectrophotometer. The remaining periphyton material from each sample was used for algal

biomass determination, expressed as grams ash-free dry mass (AFDM) per m^2 . The material was dried at 50°C for 24 h, weighed on a Sauter balance (Model AR 1014), ashed at 550°C for a minimum of 3 h, rehydrated, redried at 50°C , then cooled to ambient temperature in a desiccator and reweighed. The difference in weights equaled the AFDM of the sample.

Diatom samples were collected in each stream, after Robinson and Rushforth (1987), from three to five rock substrata representing the predominant habitat type (typically riffles). Samples were composited, preserved with 5% formalin, and returned to the laboratory. The composite sample was boiled in concentrated nitric acid, rinsed, mounted in Naphrax mountant, and examined under 1000X oil immersion using a Zeis RA microscope with Nomarski optics (St. Clair and Rushforth 1976). Relative abundances of diatom taxa were determined by counting a minimum of 1000 diatom valves from each stream. Diatoms were analyzed in terms of species richness, Simpson's index, and Kendall's coefficient of concordance (an index of similarity using all taxa with a relative abundance $>1\%$). Other algal groups such as Chlorophyta (green algae) and Cyanobacteria (blue-green algae) were not abundant at the time of sampling and thus were not considered in the present study.

One benthic sample was collected from a riffle/run habitat (pools were rare and not sampled) near each transect and analyzed for macroinvertebrates and benthic organic matter (BOM). Samples were collected using a Surber sampler ($250\ \mu\text{m}$ mesh), preserved with 5% formalin, and returned to the laboratory. Woody debris $>5\ \text{cm}$ in length that was collected in the benthic samples was rinsed of invertebrates and removed from the samples. In the laboratory, macroinvertebrates were hand-sorted from the benthic detritus with the aid of a 3X dissecting microscope, identified to the lowest feasible level (usually genus), enumerated, dried at 50°C for a minimum of 48 h, then cooled to ambient temperature in a desiccator and weighed. Dry weights, in milligrams, were determined on a Cahn (Model 25) electrobalance. The benthic detritus from each sample was used for BOM determination. The quantity of BOM, expressed as $\text{g AFDM}/\text{m}^2$, was determined as described above for periphyton. Macroinvertebrates were analyzed

in terms of density ($\text{no.}/\text{m}^2$), biomass (mg/m^2), taxa richness, Simpson's index, and relative abundances.

Chi-squared analysis was used to test for statistical differences in median substratum size between the two streams (Zar 1984). Independent sample *t* tests were used to compare the other characteristics for differences between the two streams. Prior to the *t*-test analysis all data were $\log(x + 1)$ transformed, except substratum embeddedness and the relative abundance of invertebrate taxa (both percentage measures), which were arcsine (square root $[x]$) transformed (Zar 1984). Tabular results are presented as untransformed means and standard deviations. All statistical analyses were performed on SYSTAT (Wilkinson 1990).

RESULTS

Baseflow discharge was equal in the two streams ($0.2\ \text{m}^3/\text{s}$), reflecting the similar catchment size of the burned ($22\ \text{km}^2$) and unburned ($26\ \text{km}^2$) streams. Mean baseflow width, near-bed water velocity, and BOM were not significantly different between the two streams ($P > .05$). Substratum embeddedness was significantly greater in the burned stream ($P = .01$), although the difference between mean values was not large (burned = 62.9, unburned = 52.8). It is not known whether this statistical difference was biologically meaningful or simply a reflection of the large sample size ($n = 100$).

Water depth at baseflow ($P < .01$) was lower and stream width at bankfull discharge greater ($P = .03$) in the burned stream than the unburned stream. Although not statistically significant ($P = .06$), the ratio of stream width: depth was greater in the burned than the unburned stream (216 and 91, respectively). The general appearance of the two streams was distinctly different (Fig. 1; Minshall personal observation). Large, woody debris and streamside riparian vegetation, which provide bank and channel stability, were noticeably absent in the burned stream.

Mean substratum size was not significantly different between the two streams ($P > .05$) in 1993, possibly because large boulders were not recorded in the measurements (see Fig. 1). We collected additional data on substratum size in August 1994 and included large boulders in the measurements. Further, substrata within



Fig. 1. Representative photographs of the burned (upper) and unburned (lower) streams five years after the 1988 wildfire. Note absence of large, woody debris and streamside riparian vegetation in the burned stream.

the bankfull channel were measured in 1994, in contrast to measures being recorded only within the baseflow channel in 1993. The 1994 results showed that mean substratum size was significantly larger in the unburned than in the burned stream ($P < .01$).

A comparison of median substratum size showed similar results to that of mean substratum size. Median substratum size was not different between the two streams when measurements excluded large boulders and were confined to the baseflow channel ($P > .05$). However, when measurements included large boulders and encompassed the bankfull channel, the difference in median size was significant ($P < .01$). Whether large boulders were present in the burned stream prior to the wildfire has yet to be determined. However, in other streams influenced by intensive wildfire, large boulders were observed to be buried by inorganic debris (primarily gravel and fine sediments) within five years following wildfire (Minshall personal observation).

The burned stream contained less periphyton chlorophyll *a* ($P = .06$) and AFDM ($P < .01$) than did the unburned stream (Table 1). Diatom species richness was greater in the burned (34 taxa) than in the unburned stream (27 taxa; Table 2). Simpson's index was lower for the burned than the unburned stream (0.12 and 0.42, respectively). Kendall's coefficient of concordance for the two diatom communities was 0.22, suggesting distinct assemblage composition among sites. For example, *Navicula perimitis* Hust. was the most abundant species in the burned stream, constituting 24.7% of

the assemblage, while *Hannaea arcus* (Ehr.) Patr. comprised 63.1% of the assemblage in the unburned stream (Table 2).

Mean macroinvertebrate density and biomass were lower in the burned than unburned stream (Table 3), but the differences were not significant ($P > .05$). For example, mean density in the burned stream was 9960 individuals/m², while the unburned stream had 16,950 individuals/m², and mean biomass (dry weight) was 1960 and 3200 mg/m² in the burned and unburned streams, respectively. Taxa richness and Simpson's index both were reduced in the burned stream, although the difference was significant only for Simpson's index ($P = .04$) (Table 3). The burned stream contained a mean of 15 taxa per benthic sample compared to a mean of 20 taxa for the unburned stream. The mean Simpson's index was 0.57 for the burned stream and 0.73 for the unburned stream. Chironomidae was the most abundant taxon in both streams (Table 4), although their relative abundance was significantly greater ($P = .03$) in the unburned stream. There were no statistical differences ($P > .05$) in relative abundances of other taxa common to both streams (Hydracarina, Simuliidae, *Baetis bicaudatus*, *Cinygmula*, and *Zapada columbiana*).

DISCUSSION

Alterations of the surrounding terrestrial landscape by major unpredictable disturbances such as hurricanes, volcanic eruptions, or wildfire directly influence streams draining the

TABLE 1. Means (SD) and *P* values for physical characteristics measured in the study streams.

	Burned	Unburned	<i>P</i> value
Baseflow width (m)	5.9 (2.6)	4.5 (1.8)	.46
Near-bed velocity (cm/s)	10.3 (0.1)	10.7 (0.1)	.71
BOM (g/m ²)	1.5 (0.8)	2.5 (1.2)	.20
Embeddedness (%)	62.9 (28.5)	52.8 (30.0)	.01
Baseflow depth (cm)	16.9 (11.0)	24.3 (12.5)	<.01
Bankfull width (m)	35.0 (11.9)	16.6 (1.4)	.03
Bankfull width:depth ratio	216 (101)	91 (21)	.06
Periphyton chl <i>a</i> (mg/m ²)	8.9 (5.5)	32.1 (19.5)	.06
Periphyton AFDM (g/m ²)	29.2 (3.7)	50.5 (9.3)	<.01
* Mean substratum size (cm)	14.4 (10.5)	15.0 (14.2)	.26
** Mean substratum size (cm)	15.5 (15.5)	27.6 (27.4)	<.01
* Median substratum size (cm)	12.0	11.0	.72
** Median substratum size (cm)	11.0	17.0	<.01

* Baseflow channel, large boulders excluded.
 ** Bankfull channel, large boulders included.

TABLE 2. Community measures and relative abundances (%) for the diatom assemblage of each study stream.

	Burned	Unburned
Species richness	34	27
Simpson's index (C)	0.12	0.42
<i>Navicula permissis</i> Hust.	24.7	5.1
<i>Nitzschia dissipata</i> (Kuetz.) Grun.	17.3	4.2
<i>Achnanthes lanceolata</i> (Breb.) Grun.	9.8	1.7
<i>Nitzschia paleacea</i> Grun.	7.6	3.1
<i>Navicula arvensis</i> Hust.	4.8	2.4
<i>Hannaea arcus</i> (Ehr.) Patr.	2.1	63.1

affected watersheds. For example, the Mt. St. Helens eruption of 1980 dramatically changed drainage patterns and river networks, eliminated terrestrial vegetation, and caused major debris flows that scoured stream channels (Wilzbach et al. 1983, Hawkins 1988). However, high spatial variation in the intensity of these major disturbances may occur, causing temporal differences in recovery patterns (Yount and Niemi 1990). In catchments of YNP the relative area burned ranged from <10% to >90% (Minshall and Brock 1991). Further, the degree of alteration of stream habitat was highly correlated with percent of catchment burned (Robinson and Minshall in press).

In the present study significant differences were observed in the benthic habitat of the two streams. The width:depth ratio of the burned stream was greater than that of the unburned stream. Anderson (1992) also observed increased width:depth ratios following major disturbances in streams of the Cascade Mountains. With large boulders included in the measurements, the unburned stream exhibited significantly greater substratum size. Gurtz and Wallace (1984) demonstrated that large substrata could mediate the effects of large-scale disturbances by providing stable habitat for benthic organisms. At the time of sampling, the burned stream did not contain the larger-sized substrata found in the unburned stream. It is probable that the larger substrata in the burned stream were buried by inorganic sediments following the wildfire, as has been observed in other YNP streams (Minshall personal observation). Thus, one effect of the wildfire appeared to be alteration of the substrata in such a manner as to make the benthic habitat more susceptible to future disturbances (e.g., Gurtz and Wallace 1984).

TABLE 3. Means (SD) and P values of macroinvertebrate community measures for the study streams.

	Burned	Unburned	P value
Density (no./m ²)	9963 (4730)	16,948 (8998)	.31
Biomass (mg/m ²)	1956 (1056)	3198 (1274)	.19
Taxa richness	15.0 (3.7)	20.4 (2.6)	.58
Simpson's index (C)	0.57 (0.09)	0.73 (0.10)	.04

Lamberti et al. (1991) found that faunal densities and macroinvertebrate species richness had recovered within one year following a major debris flow in an Oregon stream. In central Idaho, however, streams disturbed by wildfire and unburned reference streams showed little similarity in macroinvertebrate assemblages, even after five years of recovery (Richards and Minshall 1992). Similarly, in the present study the influence of wildfire was still apparent after five years. Macroinvertebrate community structure was not similar between the two systems, despite their close proximity to each other (0.5 km). Most researchers agree that recovery of the benthic community will correspond to recovery of the surrounding landscape (Steinman and Lamberti 1988, Minshall et al. 1989, Lamberti et al. 1991, Minshall and Brock 1991, Anderson 1992, Richards and Minshall 1992, but see Hawkins 1988).

Primary producers (lotic algae) may recover sooner than consumers (macroinvertebrates and fish) because of their much shorter life cycles, and subsequently they may influence recovery of the higher trophic levels (Steinman and McIntire 1990). In the present study, periphyton biomass (as AFDM) in the unburned stream was 1.7X greater than in the burned stream, implying a present lack of recovery by primary producers in the burned system. Macroinvertebrate taxa richness also was greater in the unburned stream than in the burned stream. How functional or structural recovery of macroinvertebrates is related to algal recovery following wildfire has yet to be determined, but provides an interesting and important avenue for future research. Algae have shorter life cycles and reduced mobility relative to macroinvertebrates, and possibly the two groups respond differently to large-scale disturbances.

TABLE 4. Mean (SD) density and relative abundance of the 10 most abundant macroinvertebrate taxa from each stream. These taxa constituted >90% of the assemblage in their respective streams.

Taxa	Density (no./m ²)	Relative abundance (%)
----- Burned -----		
Chironomidae	5437 (2218)	59.7 (19.2)
Simuliidae	2737 (4583)	18.2 (24.1)
<i>Baetis bicaudatus</i>	576 (346)	8.1 (6.3)
Hydracarina	148 (232)	1.5 (2.3)
<i>Epeorus albertae</i>	95 (123)	1.0 (1.2)
<i>Zapada columbiana</i>	90 (72)	1.0 (0.7)
<i>Epeorus longimanus</i>	80 (125)	0.5 (0.7)
<i>Cinygmula</i>	75 (71)	0.8 (0.7)
<i>Rhythrogena</i>	37 (52)	0.3 (0.3)
<i>Anelctus cooki</i>	22 (15)	0.3 (0.2)
----- Unburned -----		
Chironomidae	14,676 (8289)	84.7 (6.7)
Hydracarina	372 (253)	2.1 (1.6)
<i>Cinygmula</i>	314 (187)	1.7 (0.8)
<i>Zapada columbiana</i>	310 (215)	1.6 (0.6)
Simuliidae	307 (181)	3.8 (5.3)
<i>Baetis bicaudatus</i>	125 (60)	1.4 (1.7)
<i>Drumella coloradensis</i>	75 (64)	0.4 (0.5)
<i>Rhyacophila angelita</i>	73 (55)	0.4 (0.2)
<i>Rhyacophila tucula</i>	52 (29)	0.3 (0.1)
<i>Kogotus</i>	52 (35)	0.3 (0.2)

After five years of recovery, the channel of the burned stream still appeared unstable as indicated by different diatom assemblages between the two streams. For example, the small, adnate diatom *Navicula permitis* Hust. was predominant in the burned stream but was found in relatively low abundance in the unburned stream. *N. permitis* was predominant in other YNP streams influenced by the 1988 wildfires, and it has been suggested that a diatom community with an abundance of *N. permitis* is indicative of more physically disturbed stream environments (Robinson et al. 1994). Further, Robinson et al. (1994) showed diatom recovery among 14 streams in Yellowstone was inversely related to degree of disturbance by wildfire. Similarly, Steinman and Lamberti (1988) found little recovery, after six years, in the composition of algal communities in intensively disturbed streams of Mt. St. Helens. In summary, benthic community recovery patterns appeared to be related to the recovery of stream physical habitat which, five years after the 1988 wildfires, still displayed evidence of instability. When examining the recovery of benthic communities following

large-scale disturbance, one must remain aware of the connections between the terrestrial landscape, lotic habitat, and benthic organisms.

ACKNOWLEDGMENTS

We thank Vincent Archer, Michael Bray, Justin Gill, and especially Scott Relyea for assistance in the field. Cecily Nelson, Jason Nelson, Mark Overfield, and Jeffrey Varricchione assisted in the laboratory. Suggestions from Dr. Richard Hauer and two anonymous reviewers greatly improved the manuscript. The research was partially supported by Grant No. 725 from the Faculty Research Committee, Idaho State University. Additional funding was provided through Yellowstone Ecosystem Studies (Dr. Robert Crabtree, Director) and the Department of Botany and Range Sciences, Brigham Young University.

LITERATURE CITED

ANDERSON, N. H. 1992. Influence of disturbance on insect communities in Pacific Northwest streams. *Hydrobiologia* 248: 79–92.

GURTZ, M. E., AND J. B. WALLACE. 1984. Substrate-mediated response of stream invertebrates to disturbance. *Ecology* 65: 1556–1569.

HAWKINS, C. P. 1988. Effects of watershed vegetation and disturbance on invertebrate community structure in western Cascade streams: implications for stream ecosystem theory. *Verhandlungen der Internationale Vereinigung für Theoretische und Angewandte Limnologie* 23: 1167–1173.

HOLM-HANSEN, O., AND B. RIEMANN. 1973. Chlorophyll *a* determination: improvements in methodology. *Oikos* 30: 438–447.

HURLBERT, S. H. 1984. Pseudoreplication and the design of ecological field experiments. *Ecological Monographs* 54: 187–211.

LAMBERTI, G. A., S. V. GREGORY, L. R. ASHENAS, R. C. WILDMAN, AND K. M. S. MOORE. 1991. Stream ecosystem recovery following a catastrophic debris flow. *Canadian Journal of Fisheries and Aquatic Sciences* 48: 196–208.

MCAULIFFE, J. R. 1984. Competition for space, disturbance, and the structure of a benthic stream community. *Ecology* 65: 894–908.

MIHUC, T., G. W. MINSHALL, AND C. T. ROBINSON. In press. Responses of benthic macroinvertebrate populations in Cache Creek Yellowstone National Park to the 1988 wildfire. In: D. G. Despain and P. Schullery, editors, *The ecological implications of fire in Greater Yellowstone: 2nd biennial conference on the Greater Yellowstone Ecosystem*. National Park Service, Yellowstone National Park.

MINSHALL, G. W. 1988. Stream ecosystem theory: a global perspective. *Journal of the North American Benthological Society* 7: 263–288.

- MINSHALL, G. W., AND J. T. BROCK. 1991. Observed and anticipated effects of forest fire on Yellowstone stream ecosystems. Pages 123–135 in R. B. Keiter and M. S. Boyce, editors, *The Greater Yellowstone Ecosystem: redefining America's wilderness heritage*. Yale University Press, New Haven, CT.
- MINSHALL, G. W., AND R. C. PETERSEN. 1985. Towards a theory of macroinvertebrate community structure in stream ecosystems. *Archives für Hydrobiologia* 104: 49–76.
- MINSHALL, G. W., J. T. BROCK, AND J. D. VARLEY. 1989. Wild-fire and Yellowstone's stream ecosystems. *BioScience* 39: 707–715.
- O'NEILL, R. V., D. L. DEANGELIS, J. B. WAIDE, AND T. H. F. ALLEN. 1986. A hierarchical concept of ecosystems. Princeton University Press, Princeton, NJ. 257 pp.
- PLATTS, W. S., W. F. MEGAHAN, AND G. W. MINSHALL. 1983. Methods for evaluating stream, riparian, and biotic conditions. USDA Intermountain Forest and Range Experiment Station, Ogden, UT. General Technical Report INT-138. 70 pp.
- RESH, V. H., ET AL. 1988. The role of disturbance in stream ecology. *Journal of the North American Benthological Society* 7: 433–455.
- RICHARDS, C., AND G. W. MINSHALL. 1992. Spatial and temporal trends in stream macroinvertebrate communities: the influence of catchment disturbance. *Hydrobiologia* 241: 173–184.
- ROBINSON, C. T., AND G. W. MINSHALL. 1986. Effects of disturbance frequency on stream benthic community structure in relation to canopy cover and season. *Journal of the North American Benthological Society* 5: 237–248.
- . In press. Physical and chemical responses of streams in Yellowstone following the 1988 wildfire. In D. G. Despain and P. Schullery, editors, *The ecological implications of fire in Greater Yellowstone: 2nd biennial conference on the Greater Yellowstone Ecosystem*. National Park Service, Yellowstone National Park.
- ROBINSON, C. T., AND S. R. RUSHFORTH. 1987. Effects of physical disturbance and canopy cover on attached diatom community structure in an Idaho stream. *Hydrobiologia* 154: 49–59.
- ROBINSON, C. T., G. W. MINSHALL, AND L. VAN EVERY. 1993. Seasonal trends and colonization patterns of macroinvertebrate assemblages in two streams with contrasting flow regimes. *Great Basin Naturalist* 53: 321–331.
- ROBINSON, C. T., S. R. RUSHFORTH, AND G. W. MINSHALL. 1994. Diatom assemblages of streams influenced by wildfire. *Journal of Phycology* 30: 209–216.
- SCARSBROOK, M. R., AND C. L. TOWNSEND. 1993. Stream community structure in relation to spatial and temporal variation: a habitat templet study of two contrasting New Zealand streams. *Freshwater Biology* 29: 395–410.
- ST. CLAIR, L. L., AND S. R. RUSHFORTH. 1976. The diatom flora of the Goshen Warm Springs Ponds and Wet Meadows, Goshen, Utah, U.S.A. *Nova Hedwigia* 24: 353–425.
- STEINMAN, A. D., AND G. A. LAMBERTI. 1988. Lotic algal communities in the Mt. St. Helens region six years following the eruption. *Journal of Phycology* 24: 482–489.
- STEINMAN, A. D., AND C. D. MCINTIRE. 1990. Recovery of lotic periphyton communities. *Environmental Management* 14: 589–604.
- STRAHLER, A. N. 1952. Hypsometric (area-altitude) analysis of erosional topography. *Geological Society of America Bulletin* 63: 1117–1142.
- TOWNSEND, C. R. 1989. The patch dynamics concept of stream community ecology. *Journal of the North American Benthological Society* 8: 36–50.
- WALLACE, J. B., D. S. VOGEL, AND T. F. CUFFNEY. 1986. Recovery of a headwater stream from an insecticide-induced community disturbance. *Journal of the North American Benthological Society* 5: 115–126.
- WILKINSON, L. 1990. SYSTAT: the system for statistics. SYSTAT, Inc., Evanston, IL. 677 pp.
- WILZBACH, M. A., T. H. DUDLEY, AND J. D. HALL. 1983. Recovery patterns in stream communities impacted by the Mt. St. Helens eruption. *Water Resources Research Institute, Oregon State University, Corvallis*. Report No. WRR1-83. 33 pp.
- YOUNT, J. D., AND G. J. NIEMI. 1990. Recovery of lotic communities and ecosystems from disturbance—a narrative review of case studies. *Environmental Management* 14: 547–569.
- ZAR, J. H. 1984. *Biostatistical analysis*. Prentice-Hall Inc., Englewood Cliffs, NJ. 718 pp.

Received 16 November 1994

Accepted 2 March 1995